



A continuous publication, open access, peer-reviewed journal

ACCESS ONLINE

CORRIGENDUM

Corrigendum: Breakthroughs in hereditary angioedema management: a systematic review of approved drugs and those under research

Abstract

The authors wish to make corrections to their article:

Nicola S, Rolla G, Brussino L. Breakthroughs in hereditary angioedema management: a systematic review of approved drugs and those under research. Drugs in Context 2019; 8: 212605. DOI: 10.7573/dic.212605

Corrigendum

The authors regret that there are errors in their original paper.

1. The following shows where original text has been removed (strikethrough) and new wording introduced (underlined).

Page 3, section on Berinert®

In 2009, the <u>US</u> FDA approved Berinert[®] (CSL Behring), a pasteurized plasma-derived C1-INH concentrate, available for intravenous (IV) administration in patients needing on-demand therapy. This drug was <u>also not</u> approved for short-term prophylaxis <u>in the EU in 2013</u>. but is used off-label for this-purpose.

The following reference from the Head of Medicine website (EU approval document) should also be added to this statement:

https://mri.cts-mrp.eu/Human/Downloads/DE_H_0481_003_ PAR.pdf

Page 5, section on Haegarda®

C1-INH (SC) Haegarda®

Haegarda[®] and Berinert[®] 2000/3000 (CSL Behring) were approved (US and EU) was approved by the FDA in 2017 as the first subcutaneous C1 inhibitor concentrate, also indicated for self-administration, at the dose of 60 IU/kg twice weekly. Nevertheless, the European Medicines Agency did not formallyapprove the marketing of the drug.²⁹

 Table 1 lists two products Berinert[®] and Haegarda[®] on separate lines; however, they should be grouped together as follows:

Drug name	Active substance	Type and action	Administration	Indication	T1/2
Berinert®, CSL Behring	C1-INH	Plasma-derived C1-INH concentrate pasteurized, nanofiltered	IV	On-demand, short-term prophylaxis	36-48 hours
Berinert® 2000/3000 (Haegarda® brand name in the United States and Canada)	C1-INH	Plasma-derived C1-INH concentrate pasteurized, nanofiltered	SC	Long-term prophylaxis	69 hours